

# ANDROCUR<sup>®</sup>-100 Fact Sheet

*cyproterone acetate*

Information in this fact sheet is meant to assist you in making decisions about your treatment. Always make medication decisions in consultation with your health team.

Name:	ANDROCUR <sup>®</sup> -100 ( <i>cyproterone acetate</i> )
What is Androcur <sup>®</sup> -100 used for?	<p>Androcur<sup>®</sup>-100 is a steroidal anti-androgen. The active ingredient is cyproterone acetate.</p> <p>Androgens such as testosterone are natural male sex hormones. In some types of prostate cancer, androgens may help the cancer cells to grow. By blocking these hormones, Androcur<sup>®</sup>-100 may slow or stop the growth of cancer.</p> <p>Androcur<sup>®</sup>-100 may be used in combination with other medicines or following surgical removal of the testes to treat side effects such as “hot flushes” or “sweats” and to prevent any initial worsening of the disease.</p> <p><b>Ask your doctor if you have any questions about why this medicine has been prescribed for you.</b> This medicine is available only with a doctor’s prescription. Your doctor may have prescribed it for another reason.</p>
What does Androcur <sup>®</sup> -100 look like?	Androcur <sup>®</sup> -100 tablets are white, capsule-shaped tablets, with “LA” markings on the scored side and a hexagon marking on the other side, presented in blisters of 10. Each pack contains 50 tablets.
How is Androcur <sup>®</sup> -100 given?	<p><b>Swallow the tablets whole with some liquid after meals.</b></p> <p><b>Take your medicine after meals at about the same time each day to maximise effect and to help you remember when to take them.</b> Missed Androcur<sup>®</sup>-100 tablets may diminish the effectiveness of treatment.</p> <p><b>Do not take a double dose to make up for a dose that you have missed.</b> This may increase the chance of you getting an unwanted side effect.</p>

What are the common side effects?

**Tell your doctor, nurse or pharmacist as soon as possible if you do not feel well while you are taking Androcur®-100.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

**Do not be alarmed by the following lists of side effects.** You may not experience any of them.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor, nurse or pharmacist if you notice any of the following and they worry you:**

- tiredness, fatigue
- weight change
- headache
- depressed mood
- nausea and other gastrointestinal complaints
- decreased sexual drive
- breast pain, change in breast size, breast swelling and/or tenderness
- breast enlargement in men
- hot flushes, sweating
- shortness of breath
- osteoporosis.

If you were fertile before treatment, Androcur®-100 will normally prevent sperm production. Fertility is usually regained within a few months of discontinuing therapy.

Androcur®-100 will also normally result in the inability to get or maintain an erection (impotence).

What are the less common side effects?

**Tell your doctor, nurse or pharmacist if you notice anything that is making you feel unwell.** Other side effects not listed may happen in some people.

**Tell your doctor immediately or go to an Emergency Department at your nearest hospital if you notice any of the following:**

- yellowing of the skin and/or eyes
- light-coloured bowel motions
- dark coloured urine
- severe upper abdominal pain
- vomiting blood or material that looks like coffee grounds
- bleeding from the back passage
- black sticky bowel motions (stools) or bloody diarrhoea
- sudden severe headache
- loss of vision
- loss of coordination
- slurred speech
- shortness of breath
- chest pain
- numbness
- heat or swelling in the arms and legs.

The above list includes serious side effects that may require medical attention or hospitalisation. Serious side effects are rare.

## Reference

- TGA Consumer Medical Information (CMI) Sept 2012, Androcur®- 100, Therapeutic Goods Administration, Canberra, viewed 4<sup>th</sup> July 2013.  
<https://www.ebs.tga.gov.au/ebs/picmi/picmirepository.nsf/pdf?OpenAgent&id=CP-2010-CMI-03415-3>

## Disclaimer

Prostate Cancer Foundation of Australia develops materials based on best available evidence and takes advice from recognised experts in the field in developing such resource; however it cannot guarantee and assumes no legal responsibility for the currency or completeness of the information.